



TITLE:

Holographic Branching and Entanglement Renormalization(New Development of Numerical Simulations in Low-Dimensional Quantum Systems: From Density Matrix Renormalization Group to Tensor Network Formulations)

AUTHOR(S):

Evenbly, Glen; Vidal, Guifre

---

CITATION:

Evenbly, Glen ...[et al]. Holographic Branching and Entanglement Renormalization(New Development of Numerical Simulations in Low-Dimensional Quantum Systems: From Density Matrix Renormalization Group to Tensor Network Formulations). 物性研究 2011, 9 ...

ISSUE DATE:

2011-03-05

URL:

<http://hdl.handle.net/2433/169456>

RIGHT:

# Holographic Branching and Entanglement Renormalization

Glen Evenbly and Guifre Vidal

School of Mathematics and Physics, the University of Queensland, QLD 4072, Australia

Entanglement renormalization [1] is a coarse-graining transformation for quantum lattice systems. It produces the multi-scale entanglement renormalization ansatz (MERA) [2], a tensor network state used to represent ground states of strongly correlated systems in one and two spatial dimensions [3, 4]. In 1D, the MERA is known to reproduce the logarithmic violation of the boundary law for entanglement entropy,  $S_L \approx \log L$ , characteristic of critical ground states. In contrast, in 2D the MERA strictly obeys the entropic boundary law,  $S_L \approx L$ , characteristic of gapped systems and a class of critical systems. Therefore a number of highly entangled 2D systems, such as free fermions with a 1D Fermi surface, Fermi liquids and spin Bose metals, which display a logarithmic violation of the boundary law,  $S_L \approx L \log L$ , cannot be described by a regular 2D MERA.

It is well-known that at low energies, a many-body system may decouple into two or more independent degrees of freedom (e.g. spin-charge separation in 1D systems of electrons). In this talk I will explain how, in systems where low energy decoupling occurs, entanglement renormalization can be used to obtain an explicit decoupled description. The resulting tensor network state, the *branching MERA* [5], can reproduce a logarithmic violation of the boundary law in 2D and, as additional numeric evidence also suggests, might be a good ansatz for the highly entangled systems with a 1D Fermi (or Bose) surface mentioned above. In addition, after recalling that the MERA can be regarded as a specific (discrete) realization of the holographic principle [6], we will see that the branching MERA leads to exotic holographic geometries.

## References

- [1] G. Vidal, Phys. Rev. Lett. **99**, 220405 (2007).
- [2] G. Vidal, Phys. Rev. Lett. **101**, 110501 (2008).
- [3] G. Evenbly and G. Vidal, Phys. Rev. Lett. **102**, 180406 (2009).
- [4] G. Evenbly and G. Vidal, Phys. Rev. Lett. **104**, 187203 (2010).
- [5] G. Evenbly and G. Vidal, *in preparation*.
- [6] B. Swingle, arXiv:0905.1317v1.